

ON A QUASI-LOCAL MASS

XIAO ZHANG

ABSTRACT. We modify previous quasi-local mass definition. The new definition provides expressions of the quasi-local energy, the quasi-local linear momentum and the quasi-local mass. And they are equal to the ADM expressions at spatial infinity. Moreover, the new quasi-local energy has the positivity property.

1. INTRODUCTION

Recently, there are many attempts to define quasi-local mass [7, 8, 9, 10, 19, 16, 20, 21, 11] using certain idea initiated by Brown and York [1, 2]. The definitions are valid for spacelike 2-surfaces. The positivity property requires the 2-surfaces are spheres which are in certain initial data sets.

It is well-known that the Brown-York mass decreases for the round sphere in time slices in the Schwarzschild spacetime. In order to obtain a quasi-local mass of Brown-York's type with increasing monotonicity property, we proposed a new definition by choosing certain spinor norm as lapse function [23, 24]. But, as pointed out in [23], our quasi-local mass are not equal to the ADM mass at spatial infinity.

In this note, we modify our definition and provide expressions of the quasi-local energy, the quasi-local linear momentum and the quasi-local mass. In particular, we observe that the norm of quasi-local linear momentum 3-vector is well-defined although this 3-vector can not be defined in a quasi-local sense. This is unlike the ADM total linear momentum at spatial infinity, which both the norm and the 3-vector are well-defined. We show also that these quantities are equal to the corresponding ADM quantities at spatial infinity. Moreover, the new modified quasi-local energy has the positivity property. In particular, the vanishing of the quasi-local mass implies the spacetime is flat along the 2-spheres, which is in consequence of applying Brown and York's isometric embedding approach. This later result does not seem to hold true in the other recent definitions of quasi-local mass mentioned above.

The essential idea of our approach in this note as well as in [23, 24] is to localize Witten's proof of the positive energy theorem [14, 15, 22, 12] to the (spacelike) bounded domain. The similar approach was used by Dougan and Mason earlier to define the quasi-local mass [4]. In

our previous formulation, the 2-spheres are assumed to be apparent horizon only. In order that the Dirac-Witten equation has a solution, we have to assume that the mean curvature of spacelike hypersurfaces does not change sign. In this note we include the case that at least one boundary 2-sphere satisfies the strict inequality in apparent horizon conditions. This will ensure the existence of the Dirac-Witten equation on any spacelike hypersurfaces whose mean curvature may change sign, and this existence theorem should be required in Dougan and Mason's definition [4, 17, 18].

Surprisingly, our quasi-local mass increases for the round sphere in time slices in the Schwarzschild spacetime [23] although we are not able to prove the increasing monotonicity property in general case. This should have deep physical reason which can be seen naively as follows: In classical mechanics, the total mass is the space integral of the local mass density. But this fact is no longer true in general relativity as vacuum can have the total energy and the total mass as well. However, Witten's energy formula [22, 12] indicates that the total energy and the total mass are the space integrals of the "generalized" local mass density involving both the energy-momentum tensors and spinors. From this point of view, we think our approach is in a way toward the full understanding of quasi-local quantities.

2. DIRAC-WITTEN EQUATIONS

Let $(N, \tilde{g})$ be a 4-dimensional spacetime which satisfies the Einstein fields equations. Let (M, g, p) be a smooth initial data set. Fix a point $z \in M$ and an orthonormal basis $\{e_\alpha\}$ of $T_z N$ with e_0 future-time-directed normal to M and e_i tangent to M ($1 \leq i \leq 3$).

Denote by $\mathbb{S}$ the (local) spinor bundle of N . It exists globally over M and is called the hypersurface spinor bundle of M . Let $\tilde{\nabla}$ and $\bar{\nabla}$ be the Levi-Civita connections of $\tilde{g}$ and g respectively, the same symbols are used to denote their lifts to the hypersurface spinor bundle. There exists a Hermitian inner product $(\cdot, \cdot)$ on $\mathbb{S}$ along M which is compatible with the spin connection $\tilde{\nabla}$. The Clifford multiplication of any vector $\tilde{X}$ of N is symmetric with respect to this inner product. However, this inner product is not positive definite. As $e_0^2 = \text{id}$, $\mathbb{S}$ is decomposed into the direct sums of two subbundles which are the eigenbundles of e_0 along M . This indicates there exists a positive definite Hermitian inner product defined by $\langle \cdot, \cdot \rangle = (e_0 \cdot, \cdot)$ on $\mathbb{S}$ along M .

The second fundamental form of the initial data set is defined as $p_{ij} = \tilde{g}(\tilde{\nabla}_i e_0, e_j)$. Suppose M has boundary Σ which has finitely many connected components $\Sigma^1, \dots, \Sigma^l$, each of which is a topological 2-sphere and endowed with its induced Riemannian and spin structures. Fix a point $z \in \Sigma$ and an orthonormal basis $\{e_i\}$ of $T_z M$ with $e_r = e_1$

outward normal to Σ and e_a tangent to Σ for $2 \leq a \leq 3$. Let $h_{ab} = \langle \bar{\nabla}_a e_r, e_b \rangle$ be the second fundamental form of Σ . Let $H = \text{tr}(h)$ be its mean curvature. Σ is a *future/past apparent horizon* if

$$H \mp \text{tr}(p|_\Sigma) \geq 0 \quad (2.1)$$

holds on Σ .

Denote by ∇ the lift of the Levi-Civita connection of Σ to the spinor bundle $\mathbb{S}|_\Sigma$. The Dirac-Witten operator along M , the Dirac operator of M acting on $\mathbb{S}$ and the Dirac operator of Σ acting on $\mathbb{S}|_\Sigma$ are defined as $\tilde{D} = e_i \cdot \tilde{\nabla}_i$, $\bar{D} = e_i \cdot \bar{\nabla}_i$, $D = e_a \cdot \nabla_a$ respectively. Let $P_\pm = \frac{1}{2}(Id \pm e_0 \cdot e_r \cdot)$ be the projective operators on $\mathbb{S}|_\Sigma$. Then $\mathbb{S}|_\Sigma \cong \mathbb{S}|_\Sigma^+ \oplus \mathbb{S}|_\Sigma^-$ is decomposed into the direct sum of two equivalent complex 2-dimensional sub-bundles $\mathbb{S}|_\Sigma^\pm = \{\phi \in \mathbb{S}|_\Sigma : P_\pm \phi = \phi\}$.

The following integral identity is given by the relation $\tilde{\nabla}_a = \nabla_a + \frac{1}{2}h_{ab}e_r \cdot e_b \cdot - \frac{1}{2}p_{aj}e_0 \cdot e_j \cdot$ and the Weitzenböck type formula for the Dirac-Witten operator

$$\begin{aligned} \int_M |\tilde{\nabla}\phi|^2 + \langle \phi, \mathcal{T}\phi \rangle - |\tilde{D}\phi|^2 &= \int_\Sigma \langle \phi, (e_r \cdot D - \frac{H}{2})\phi \rangle \\ &+ \frac{1}{2} \int_\Sigma \langle \phi, \text{tr}(p|_\Sigma)e_0 \cdot e_r \cdot \phi \rangle - \frac{1}{2} \int_\Sigma \langle \phi, p_{ar}e_0 \cdot e_a \cdot \phi \rangle \end{aligned} \quad (2.2)$$

where $\mathcal{T} = \frac{1}{2}(T_{00} + T_{0i}e_0 \cdot e_i \cdot)$. Let $\phi = \phi^+ + \phi^-$. Since $e_0 \cdot e_r \cdot \phi^\pm = \pm \phi^\pm$, we have $e_0 \cdot e_r \cdot (e_r \cdot D\phi^\pm) = \mp e_r \cdot D\phi^\pm$ and $e_0 \cdot e_r \cdot (e_0 \cdot e_a \cdot \phi^\pm) = \mp e_0 \cdot e_a \cdot \phi^\pm$, therefore the right hand side of (2.2) is

$$\begin{aligned} \text{RHS} &= \int_\Sigma \langle \phi^+, e_r \cdot D\phi^- \rangle + \langle \phi^-, e_r \cdot D\phi^+ \rangle \\ &- \frac{1}{2} \int_\Sigma [H - \text{tr}(p|_\Sigma)] |\phi^+|^2 - \frac{1}{2} \int_\Sigma [H + \text{tr}(p|_\Sigma)] |\phi^-|^2 \\ &- \frac{1}{2} \int_\Sigma \langle \phi^+, p_{ar}e_0 \cdot e_a \cdot \phi^- \rangle - \frac{1}{2} \int_\Sigma \langle \phi^-, p_{ar}e_0 \cdot e_a \cdot \phi^+ \rangle. \end{aligned}$$

Parts (a) and (b) of the following existences are proved in [23] which correspond to the case all Σ_i are either future or past apparent horizons. Part (c) corresponds to the case that at least one boundary 2-sphere satisfies the strict inequality in apparent horizon conditions.

Proposition 1. *Let $(N, \tilde{g})$ be a spacetime which satisfies the dominant energy condition. Let (M, g, p) be a smooth initial data with the boundary Σ which has finitely many multi-components Σ_i , each of which is topological 2-sphere.*

(a) If $\text{tr}_g(p) \geq 0$ and all Σ_i are past apparent horizons, then the following Dirac-Witten equation has a unique smooth solution $\phi \in \Gamma(\mathbb{S})$

$$\begin{cases} \tilde{D}\phi &= 0 & \text{in } M \\ P_+\phi &= P_+\phi_{0i} & \text{on } \Sigma_i \end{cases} \quad (2.3)$$

for given $\phi_{0i} \in \Gamma(\mathbb{S}|_{\Sigma_i})$;

(b) If $\text{tr}_g(p) \leq 0$ and all Σ_i are future apparent horizons, then the following Dirac-Witten equation has a unique smooth solution $\phi \in \Gamma(\mathbb{S})$

$$\begin{cases} \tilde{D}\phi &= 0 & \text{in } M \\ P_-\phi &= P_-\phi_{0i} & \text{on } \Sigma_i \end{cases} \quad (2.4)$$

for given $\phi_{0i} \in \Gamma(\mathbb{S}|_{\Sigma_i})$;

(c) If Σ_i , $1 \leq i \leq k_0$, are past apparent horizons and Σ_j , $k_0+1 \leq j \leq l$, are future apparent horizons, and there exists at least l_0 , $1 \leq l_0 \leq l$, such that the strict inequality in (2.1) holds for l_0 , then the following Dirac-Witten equation has a unique smooth solution $\phi \in \Gamma(\mathbb{S})$

$$\begin{cases} \tilde{D}\phi &= 0 & \text{in } M \\ P_+\phi &= P_+\phi_{0i} & \text{on } \Sigma_i \\ P_-\phi &= P_-\phi_{0j} & \text{on } \Sigma_j \end{cases} \quad (2.5)$$

for given $\phi_{0i} \in \Gamma(\mathbb{S}|_{\Sigma_i})$, $\phi_{0j} \in \Gamma(\mathbb{S}|_{\Sigma_j})$.

Proof : To establish the existences of (2.3), (2.4) and (2.5), we only need to show that any solution $\tilde{D}\phi = 0$ is trivial when $\phi_{0i} = \phi_{0j} = 0$. We first study the case (c), in this case (2.2) gives

$$\begin{aligned} \int_M |\tilde{\nabla}\phi|^2 + \langle \phi, \mathcal{T}\phi \rangle &= -\frac{1}{2} \sum_{1 \leq i \leq k_0} \int_{\Sigma_i} [H + \text{tr}(p|_{\Sigma_i})] |\phi^-|^2 \\ &\quad - \frac{1}{2} \sum_{k_0+1 \leq j \leq l} \int_{\Sigma_j} [H - \text{tr}(p|_{\Sigma_j})] |\phi^+|^2. \end{aligned}$$

Since the strict inequality in (2.1) holds for l_0 , we must have $\phi|_{\Sigma_{l_0}} = 0$ and $\tilde{\nabla}_i\phi = 0$ along M . However, in cases (a) and (b), we can not conclude that $\phi|_{\Sigma_i} = 0$ on some i directly from the apparent horizon condition (2.1). We need to use some other argument. Since $\tilde{D}(e_0 \cdot \phi) = -\text{tr}_g(p)\phi$, we have

$$\sum_{1 \leq i \leq l} \int_{\Sigma_i} \langle e_r \cdot \phi, e_0 \cdot \phi \rangle = \int_M \langle \tilde{D}\phi, e_0 \cdot \phi \rangle - \langle \phi, \tilde{D}(e_0 \cdot \phi) \rangle = \int_M \text{tr}_g(p) |\phi|^2.$$

If all Σ_i are past apparent horizons and $tr_g(p) \geq 0$, or all Σ_i are future apparent horizons and $tr_g(p) \leq 0$, then the above equality gives that $\phi|_{\Sigma_i} = 0$ for all i . We have also $\tilde{\nabla}_i \phi = 0$ along M in these cases. Now $|d|\phi|^2| \leq 2|\bar{\nabla}\phi||\phi| \leq |p||\phi|^2 \leq C_p|\phi|^2$, where $C_p = \max|p|$. If there exists a point $x_0 \in M$ such that $\phi(x_0) \neq 0$, then

$$|\phi|(x) \geq |\phi|(x_0)e^{-C_p \rho_{x_0}(x)}.$$

That taking x to the boundary Σ_{l_0} will give rise to a contradiction. Therefore $\phi = 0$ over M and the existences of (2.3), (2.4) and (2.5) follow. Q.E.D.

Proposition 2. *Let $(N, \tilde{g})$ be a spacetime which satisfies the dominant energy condition. Let (M, g, p) be a smooth initial data with the boundary Σ which has finitely many multi-components Σ_i , each of which is topological 2-sphere. Suppose ϕ, ψ are smooth spinors which satisfy $\tilde{\nabla}_i \phi = 0, \tilde{\nabla}_i \psi = 0$. Denote $\phi_0^\pm = P_\pm \phi|_{\Sigma_i}, \psi_0^\pm = P_\pm \psi|_{\Sigma_i}$. If either ϕ_0^+, ψ_0^+ or ϕ_0^-, ψ_0^- are linearly independent on Σ_i , then ϕ, ψ are linearly independent in M .*

Proof : Let $C_1\phi(x_0) + C_2\psi(x_0) = 0$ for complex constants C_1, C_2 at some point $x_0 \in M$. We shall show that C_1, C_2 must be zero. Denote $\Phi(x) = C_1\phi(x) + C_2\psi(x)$. Let Ω be an open subset of M where Φ is nonzero. Since $\tilde{\nabla}_i \Phi = 0$, we know that $|\Phi(y)| \geq |\Phi(x)|e^{-C_p \rho(x,y)}$ for any $x, y \in \Omega$ where $C_p = \max|p|$. If Ω is nonempty, we can take y to the boundary of Ω or to the point x_0 . This will give rise to a contradiction. Therefore $\Phi(x) \equiv 0$. Hence $C_1\phi_0^\pm + C_2\psi_0^\pm = 0$ on Σ_i . Since ϕ_0^+, ψ_0^+ or ϕ_0^-, ψ_0^- is linearly independent on Σ_i , we have $C_1 = C_2 = 0$ and the proposition follows. Q.E.D.

3. QUASI-LOCAL QUANTITIES

Let $(N, \tilde{g})$ be a spacetime which satisfies the dominant energy condition. Let (M, g, p) be a smooth initial data set with boundary Σ which has finitely many connected components $\Sigma_1, \dots, \Sigma_l$, each of which is a topological 2-sphere. Suppose that some Σ_i can be smoothly isometrically embedded into a smooth spacelike hypersurface $\check{M}^3$ in the Minkowski spacetime $\mathbb{R}^{3,1}$ and denote by $\check{\aleph}$ the isometric embedding. Let $\check{\Sigma}_i$ be the image of Σ_i under the map $\check{\aleph}$. Let $\check{e}_r$ the unit vector outward normal to $\check{\Sigma}_i$ and $\check{h}_{ij}, \check{H}$ are the second fundamental form, the mean curvature of $\check{\Sigma}_i$ respectively. Denote by $p_0 = \check{p} \circ \check{\aleph}$, $H_0 = \check{H} \circ \check{\aleph}$ the pullbacks to Σ_i .

The isometric embedding $\aleph$ also induces an isometry between the (intrinsic) spinor bundles of Σ_i and $\check{\Sigma}_i$ together with their Dirac operators which are isomorphic to $e_r \cdot D$ and $\check{e}_r \cdot \check{D}$ respectively. This isometry can be extended to an isometry over the complex 2-dimensional sub-bundles of their hypersurface spinor bundles. Denote by $\check{\mathbb{S}}^{\check{\Sigma}_i}$ this sub-bundle of $\check{\mathbb{S}}|_{\check{\Sigma}_i}$. We choose the same orientations for N and $\mathbb{R}^{3,1}$. Since $e_2 \cdot e_3 = \check{e}_2 \cdot \check{e}_3$, $P_{\pm}$ is preserved by $\aleph$. Therefore,

$$\mathbb{S}|_{\Sigma_i}^{\pm} \cong \check{\mathbb{S}}|_{\check{\Sigma}_i}^{\pm} \cong \check{\mathbb{S}}^{\check{\Sigma}_i}. \quad (3.1)$$

Let $\check{\phi}$ be a constant section of $\check{\mathbb{S}}^{\check{\Sigma}_i}$ and denote $\phi_0 = \check{\phi} \circ \aleph$. Denote by $\check{\Xi}$ the set of all these constant spinors $\check{\phi}$ with the unit norm. This set is isometric to S^3 .

In this note, we introduce the following conditions on M :

- (i) $tr_g(p) \geq 0$, $H|_{\Sigma_i} + tr(p|_{\Sigma_i}) \geq 0$ for all i ;
- (ii) $tr_g(p) \leq 0$, $H|_{\Sigma_i} - tr(p|_{\Sigma_i}) \geq 0$ for all i ;
- (iii) $H|_{\Sigma_i} + tr(p|_{\Sigma_i}) \geq 0$ for $1 \leq i \leq k_0$, $H|_{\Sigma_i} - tr(p|_{\Sigma_i}) \geq 0$ for $k_0 + 1 \leq j \leq l$, and there exists at least l_0 , $1 \leq l_0 \leq l$, such that the strict inequality holds for l_0 .

Now we provide the definition of quasi-local quantities for some Σ_i under these conditions. Recall [23, 24]

Case 1. If Σ_i has positive Gauss curvature, then it can be embedded smoothly isometrically into $\mathbb{R}^3$ in the Minkowski spacetime $\mathbb{R}^{3,1}$. Denote by $\check{\Sigma}_i$ its image. In this case, $\check{p} = 0$.

Let ϕ be the unique solution of one of (2.3), (2.4) and (2.5) for some $\check{\phi}_i \in \check{\Xi}$ and $\check{\phi}_j = 0$ for $j \neq i$. Denote

$$\begin{aligned} m(\Sigma_i, \check{\phi}) = & \frac{area(\Sigma_i)}{8\pi \int_{\Sigma_i} |\phi|^2} \Re \int_{\Sigma_i} \left[(H_0 - H) |\phi|^2 \right. \\ & \left. + tr(p|_{\Sigma_i}) \langle \phi, e_0 \cdot e_r \cdot \phi \rangle - p_{ar} \langle \phi, e_0 \cdot e_a \cdot \phi \rangle \right]. \end{aligned}$$

Define

$$m_i^b = \min_{\check{\Xi}} m(\Sigma_i, \check{\phi}), \quad m_i^h = \max_{\check{\Xi}} m(\Sigma_i, \check{\phi}).$$

Case 2. If Σ_i has negative Gauss curvature $K_{\Sigma_i} \geq -\kappa^2$ ($\kappa > 0$) where $-\kappa^2$ is the minimum of the Gauss curvature. (Here we must choose the minimum of the Gauss curvature instead of arbitrary lower bound, otherwise the quasi-local mass defined in the following way

might depend on this arbitrary lower bound.) By [13, 3], Σ_i can be smoothly isometrically embedded into the hyperbolic space $\mathbb{H}_{-\kappa^2}^3$ with constant curvature $-\kappa^2$ as a convex surface which bounds a convex domain in $\mathbb{H}_{-\kappa^2}^3$. Let (t, x_1, x_2, x_3) be the spacetime coordinates of $\mathbb{R}^{3,1}$. Then $\mathbb{H}_{-\kappa^2}^3$ is one-fold of the spacelike hypersurfaces $\{t^2 - x_1^2 - x_2^2 - x_3^2 = \frac{1}{\kappa^2}\}$. The induced metric of $\mathbb{H}_{-\kappa^2}^3$ is $\check{g}_{\mathbb{H}_{-\kappa^2}^3} = \frac{dr^2}{1+\kappa^2 r^2} + r^2(d\theta^2 + \sin^2 \theta d\psi^2)$. It has the second fundamental form $\check{p}_{\mathbb{H}_{-\kappa^2}^3}^+ = \kappa \check{g}_{\mathbb{H}_{-\kappa^2}^3}$ for the upper-fold $\{t > 0\}$ and $\check{p}_{\mathbb{H}_{-\kappa^2}^3}^- = -\kappa \check{g}_{\mathbb{H}_{-\kappa^2}^3}$ for the lower-fold $\{t < 0\}$ with respect to the future-time-directed normal.

If condition (i) holds, we embed Σ_i into upper-fold $\{t > 0\}$. Since $\check{\Sigma}_i$ is convex, we have $\check{H} + tr(\check{p}|_{\check{\Sigma}_i}) > 0$. If condition (ii) holds, we embed Σ_i into lower-fold $\{t < 0\}$. We have $\check{H} - tr(\check{p}|_{\check{\Sigma}_i}) > 0$ in this case. If condition (iii) holds, we embed Σ_i into upper-fold if $H + tr(p|_{\Sigma_i}) \geq 0$ and embed it into lower-fold if $H - tr(p|_{\Sigma_i}) \geq 0$. Denote also by $\check{\Sigma}_i$ its image.

Let ϕ be the unique solution of one of (2.3), (2.4) and (2.5) for some $\check{\phi}_i \in \check{\Xi}$ and $\check{\phi}_j = 0$ for $j \neq i$. Denote

$$m_{\pm}(\Sigma_i, \check{\phi}) = \frac{area(\Sigma_i)}{8\pi \int_{\Sigma_i} |\phi|^2} \Re \int_{\Sigma_i} \left[(H_0 - H)|\phi|^2 + tr(p|_{\Sigma_i}) \langle \phi, e_0 \cdot e_r \cdot \phi \rangle - p_{ar} \langle \phi, e_0 \cdot e_a \cdot \phi \rangle \right] \mp \frac{area(\Sigma_i)\kappa}{4\pi \int_{\Sigma_i} |\phi|^2} \int_{\Sigma_i} \langle \phi, e_0 \cdot e_r \cdot \phi \rangle.$$

Define

$$m_i^b = \begin{cases} \min_{\check{\Xi}} m_+(\Sigma_i, \check{\phi}): & \text{if } \Sigma_i \text{ is embedded into upper-fold,} \\ \min_{\check{\Xi}} m_-(\Sigma_i, \check{\phi}): & \text{if } \Sigma_i \text{ is embedded into lower-fold.} \end{cases}$$

$$m_i^h = \begin{cases} \max_{\check{\Xi}} m_+(\Sigma_i, \check{\phi}): & \text{if } \Sigma_i \text{ is embedded into upper-fold,} \\ \max_{\check{\Xi}} m_-(\Sigma_i, \check{\phi}): & \text{if } \Sigma_i \text{ is embedded into lower-fold.} \end{cases}$$

Case 3. If Σ_i has nonnegative Gauss curvature which vanishes at some point, then it can be only $C^{1,1}$ embedded isometrically into $\mathbb{R}^3$ [5, 6]. To avoid this regularity problem, we first embed it smoothly isometrically into hyperbolic space $\mathbb{H}_{-\kappa^2}^3$ and then take the limit $\kappa \rightarrow 0$. Define

$$m_i^b = \begin{cases} \lim_{\kappa \rightarrow 0} \min_{\check{\Xi}} m_+(\Sigma_i, \check{\phi}): & \text{if } \Sigma_i \text{ is embedded into upper-fold,} \\ \lim_{\kappa \rightarrow 0} \min_{\check{\Xi}} m_-(\Sigma_i, \check{\phi}): & \text{if } \Sigma_i \text{ is embedded into lower-fold.} \end{cases}$$

$$m_i^h = \begin{cases} \lim_{\kappa \rightarrow 0} \max_{\check{\Xi}} m_+(\Sigma_i, \check{\phi}): & \text{if } \Sigma_i \text{ is embedded into upper-fold,} \\ \lim_{\kappa \rightarrow 0} \max_{\check{\Xi}} m_-(\Sigma_i, \check{\phi}): & \text{if } \Sigma_i \text{ is embedded into lower-fold.} \end{cases}$$

Remark 1. *The quasi-local quantities do not depend on the choice of spinors after taking maximum and minimum on certain constant spinor space. This procedure was not taken to rule out spinors in Dougan and Mason's definition [4].*

Remark 2. *The solutions of (2.3), (2.4) and (2.5) for some $\check{\phi}_i \in \check{\Xi}$ and $\check{\phi}_j = 0$ for $j \neq i$ are parameterized by the positive or negative part of the constant Dirac spinors in $\mathbb{R}^{3,1}$. So the set of these unit norm constant spinors is S^3 .*

Remark 3. *The normalized factor is omitted in the definitions of $m(\Sigma_i, \check{\phi})$ and $m_{\pm}(\Sigma_i, \check{\phi})$ in [23, 24].*

Definition 1. *Let $(N, \tilde{g})$ be a spacetime which satisfies the dominant energy condition. Let (M, g, p) be a smooth initial data set with the boundary Σ which has finitely many multi-components Σ_i , each of which is topological 2-sphere. If one of conditions (i), (ii), (iii) holds, then the quasi-local energy E_i^{loc} , the quasi-local linear momentum $|P_i^{loc}|$ and the quasi-local mass m_i^{loc} of Σ_i are defined respectively as follows*

$$E_i^{loc} = \frac{m_i^{\natural} + m_i^{\flat}}{2}, \quad |P_i^{loc}| = \frac{m_i^{\natural} - m_i^{\flat}}{2}, \quad m_i^{loc} = \sqrt{m_i^{\natural} m_i^{\flat}}.$$

From the following theorem, we know that $m_i^{\natural} \geq m_i^{\flat} \geq 0$. Therefore these quasi-local quantities are well-defined.

It is straightforward that these quasi-local quantities all vanish if $(N, \tilde{g})$ is the Minkowski spacetime. On the other hand, by the uniqueness, any solution of (2.3) or (2.4) with boundary value in $\check{\Xi}$ approaches a constant spinor in asymptotically flat initial data sets. Therefore, the limits at spatial infinity are

$$\lim_{r \rightarrow \infty} m_i^{\natural} = E + |P|, \quad \lim_{r \rightarrow \infty} m_i^{\flat} = E - |P|$$

where E, P_k are the ADM total energy, the ADM total linear momentum respectively and $|P| = \sqrt{P_1^2 + P_2^2 + P_3^2}$. Denote by $m = \sqrt{E^2 - P_1^2 - P_2^2 - P_3^2}$ the ADM mass. We have

$$\lim_{r \rightarrow \infty} E_i^{loc} = E, \quad \lim_{r \rightarrow \infty} |P_i^{loc}| = |P|, \quad \lim_{r \rightarrow \infty} m_i^{loc} = m.$$

Theorem 1. *Let $(N, \tilde{g})$ be a spacetime which satisfies the dominant energy condition. Let (M, g, p) be a smooth initial data set with the boundary Σ which has finitely many multi-components Σ_i , each of which is topological 2-sphere. If one of conditions (i), (ii), (iii) holds, then*

- (a) $m_i^{\flat} \geq 0 \Leftrightarrow E_i^{loc} \geq |P_i^{loc}|$ for each i ;
- (b) If $E_{i_0}^{loc} = 0$ for some i_0 , then N is flat along Σ_{i_0} .

Proof : Part (a) is a direct consequence of Proposition 1 and the following inequality proved in [23, 24]

$$\int_{\Sigma_{i_0}} \langle \phi, e_r \cdot D\phi \rangle \leq \frac{1}{2} \int_{\Sigma_{i_0}} \langle \phi, (H_0 - p_{0aa}e_0 \cdot e_r \cdot + p_{0ar}e_0 \cdot e_a \cdot) \phi \rangle. \quad (3.2)$$

For the completeness, we summarize the proof of (3.2) here. Since $\partial_i \check{\phi} = 0$, restricted on $\check{\Sigma}_{i_0}$, we have $\check{\nabla}_a \check{\phi} + \frac{1}{2} \check{h}_{ab} \check{e}_r \cdot \check{e}_b \cdot \check{\phi} - \frac{1}{2} \check{p}_{aj} \check{e}_0 \cdot \check{e}_j \cdot \check{\phi} = 0$. Pullback to Σ_{i_0} , we obtain $e_r \cdot D\phi_0^\pm = \frac{H_0}{2} \phi_0^\mp \pm \frac{1}{2} p_{0aa} \phi_0^\mp + \frac{1}{2} p_{0ar} e_0 \cdot e_a \cdot \phi_0^\pm$. Using $\int_{\Sigma_{i_0}} \langle \phi_0^-, e_r \cdot D\phi_0^+ \rangle = \int_{\Sigma_{i_0}} \langle e_r \cdot D\phi_0^-, \phi_0^+ \rangle$, we get $\int_{\Sigma_{i_0}} (H_0 - p_{0aa}) |\phi_0^+|^2 = \int_{\Sigma_{i_0}} (H_0 + p_{0aa}) |\phi_0^-|^2$.

The proof of (3.2) as well as the theorem under condition (i) or condition (iii) with Σ_{i_0} the past apparent horizon is the same as that under condition (ii) or condition (iii) with Σ_{i_0} the future apparent horizon. So we only prove the first case. Let ϕ be the smooth solution of the Dirac-Witten equation with $\phi^+ = \phi_0^+$. We have

$$\begin{aligned} \int_{\Sigma_{i_0}} \langle \phi, e_r \cdot D\phi \rangle &= 2\Re \int_{\Sigma_{i_0}} \langle \phi^-, e_r \cdot D\phi_0^+ \rangle \\ &= \Re \int_{\Sigma_{i_0}} \langle \phi^-, H_0 \phi_0^- + p_{0aa} \phi_0^- + p_{0ar} e_0 \cdot e_r \cdot \phi_0^+ \rangle \\ &\leq \frac{1}{2} \int_{\Sigma_{i_0}} (H_0 + p_{0aa}) (|\phi^-|^2 + |\phi_0^-|^2) \\ &\quad + \Re \int_{\Sigma_{i_0}} \langle \phi^-, p_{0ar} e_0 \cdot e_a \cdot \phi_0^+ \rangle \\ &= \frac{1}{2} \int_{\Sigma_{i_0}} (H_0 + p_{0aa}) |\phi^-|^2 + (H_0 - p_{0aa}) |\phi_0^+|^2 \\ &\quad + \Re \int_{\Sigma_{i_0}} \langle \phi^-, p_{0ar} e_0 \cdot e_a \cdot \phi^+ \rangle \\ &= \frac{1}{2} \int_{\Sigma_{i_0}} H_0 |\phi|^2 + p_{0aa} (|\phi^-|^2 - |\phi^+|^2) \\ &\quad + \Re \int_{\Sigma_{i_0}} \langle \phi^-, p_{0ar} e_0 \cdot e_a \cdot \phi^+ \rangle \\ &= \frac{1}{2} \int_{\Sigma_{i_0}} \langle \phi, (H_0 - p_{0aa} e_0 \cdot e_r \cdot + p_{0ar} e_0 \cdot e_a \cdot) \phi \rangle. \end{aligned}$$

For part (b), if $E_{i_0}^{loc} = 0$ for some i_0 , then $m_{i_0}^a = m_{i_0}^b = 0$. Therefore, for any $\check{\phi} \in \check{\Xi}$, $m(\Sigma_{i_0}, \check{\phi}) = 0$. This implies $\check{\nabla}_i \check{\phi} = 0$. Let $\check{\phi}_1, \check{\phi}_2 \in \check{\Xi}$ be

two linearly independent constant spinors. Let ϕ_1, ϕ_2 be the smooth solutions of one of (2.3), (2.4) and (2.5) corresponding to the boundary values $\check{\phi}, \check{\psi}$. Then $\tilde{\nabla}_i \phi_1 = \tilde{\nabla}_i \phi_2 = 0$. By Proposition 2, ϕ_1 and ϕ_2 are linearly independent. Choose the frame e_α such that $\tilde{\nabla}_\alpha e_\beta = 0$. Then

$$-\frac{1}{4}\tilde{R}_{ij\alpha\beta}e_\alpha e_\beta \phi_a = [\tilde{\nabla}_i, \tilde{\nabla}_j]\phi_a = 0 \quad (3.3)$$

for $a = 1, 2$. Note that we are not able to obtain the vanishing of $\tilde{R}_{ij\alpha\beta}$ along M as ϕ_a are complex 4-dimensional which might not belong to neither the positive eigenbundle of e_0 nor the negative eigenbundle of e_0 . However, in the proof of (3.2), the inequality $2\Re\langle\phi^-, \phi_0^-\rangle \leq |\phi^-|^2 + |\phi_0^-|^2$ is used on Σ_{i_0} , and it is forced to be the equality when $E_{i_0}^{loc} = 0$. So $\phi^- = \phi_0^-$ on Σ_{i_0} . Since the boundary value condition gives $\phi^+ = \phi_0^+$, we obtain $\phi = \phi_0$ on Σ_{i_0} . As ϕ_{0a} are pullbacks of $\check{\phi}_a$, ϕ_{0a} belong to the same eigenbundle of e_0 . This indicates the situation can be reduced to complex 2-dimensional spinor bundles on Σ_{i_0} and (3.3) holds true for two linearly independent constant spinors ϕ_{0a} on Σ_{i_0} . Therefore $\tilde{R}_{ij\alpha\beta} = 0$ on Σ_{i_0} . So $T_{00} = 0$ on Σ_{i_0} and the dominant energy condition gives $T_{\alpha\beta} = 0$ which implies $\tilde{R}_{i0\alpha\beta} = 0$ on Σ_{i_0} . Therefore the spacetime is flat along Σ_{i_0} . Q.E.D.

4. DISCUSSIONS

In this note, we define the quasi-local quantities for a 2-sphere in the spacelike hypersurface. For a spacelike 2-sphere Σ in spacetimes, we can take infimum of above defined quasi-local quantities over the set of spacelike hypersurfaces enclosed by Σ , if the set is nonempty. Because of Theorem 1, the infimum exists and is nonnegative. We can define this infimum as the quasi-local quantities of Σ . If we think the choice of the spacelike hypersurface for Σ is gauge, we can call the *infimum* gauge in this situation. Alternatively, if Σ can enclose a unique maximal hypersurface which satisfies $tr(p) = 0$ in spacetime (it is indeed the case in certain situation), then the quasi-local quantities of Σ can be defined in terms of the geometry of this 2-sphere and the maximal hypersurface. And we can call it the *maximal* gauge. The quasi-local quantities of Σ in these gauges can be understood not to depend on the spacelike hypersurface.

Acknowledgements. The author is indebted to N. Ó Murchadha, Roh-Suan Tung and Naqing Xie for some valuable conversations. This work is partially supported by NSF of China (10421001, 10725105, 10731080), NKBRPC(2006CB805905) and the Innovation Project of Chinese Academy of Sciences.

REFERENCES

- [1] J.D. Brown and J.W. York, *Quasilocal energy in general relativity*, Mathematical aspects of classical field theory (Seattle, WA, 1991), 129-142, Contemp. Math., 132, Amer. Math. Soc., Providence, RI, 1992.
- [2] J.D. Brown and J.W. York, *Quasilocal energy and conserved charges derived from the gravitational action*, Phys. Rev. D(3) 47, 1407-1419 (1993).
- [3] M.P. do Carmo, F.W. Warner, *Rigidity and convexity of hypersurfaces in spheres*, J. Diff. Geom. 4, 133-144 (1970).
- [4] A.J. Dougan and L.J. Mason, *Quasilocal mass constructions with positive energy*, Phys. Rev. Lett. 67, 2119-2122 (1991).
- [5] F. Guan, Y.Y. Li, *The Weyl problem with nonnegative Gauss curvature*, J. Diff. Geom., 39, 331-342 (1994).
- [6] J.X. Hong, C. Zuily, *Isometric embedding of the 2-sphere with nonnegative curvature in R^3* , Math. Z., 219, 323-334 (1995).
- [7] J. Kijowski, *A simple derivation of canonical structure and quasi-local Hamiltonians in general relativity*, Gen. Relat. Grav. 29, 307-343 (1997).
- [8] C-C.M. Liu and S.T. Yau, *Positivity of quasilocal mass*, Phys. Rev. Lett. 90, 231102 (2003).
- [9] C-C.M. Liu and S.T. Yau, *Positivity of quasilocal mass II*, J. Amer. Math. Soc. 19, 181 (2006).
- [10] N.Ó Murchadha, L.B. Szabados, K.P. Tod, *Comment on "Positivity of quasilocal mass"*, Phys. Rev. Lett. 92, 259001 (2004).
- [11] N.Ó Murchadha, R-S. Tung, N. Xie, *Quasi-local energy in general relativity*, arXiv: 0905.0647 [gr-qc].
- [12] T. Parker and C.H. Taubes, *On Witten's proof of the positive energy theorem*, Commun. Math. Phys. 84, 223-238 (1982).
- [13] A.V. Pogorelov, *Some results on surface theory in the large*, Adv. Math. 1, 191-264 (1964).
- [14] R. Schoen, S.T. Yau, *On the proof of the positive mass conjecture in general relativity*, Commun. Math. Phys. 65, 45-76 (1979).
- [15] R. Schoen, S.T. Yau, *Proof of the positive mass theorem. II*, Commun. Math. Phys. 79, 231-260 (1981).
- [16] Y. Shi, L-F. Tam, *Rigidity of compact manifolds and positivity of quasi-local mass*, Class. Quantum Grav. 24 2357-2366 (2007).
- [17] L.B. Szabados, *On the positivity of the quasi-local mass*, Class. Quantum Grav. 10, 1899-1905 (1993).
- [18] L.B. Szabados, *Two-dimensional Sen connections and quasi-local energy-momentum*, Class. Quantum Grav. 11, 1847-1866 (1994).
- [19] M.T. Wang, S.T. Yau, *A generalization of Liu-Yau's quasi-local mass*, Comm. Anal. Geom. 15, 249-282 (2007).
- [20] M.T. Wang, S.T. Yau, *Quasi-local mass in general relativity*, Phys. Rev. Lett. 102, 021101 (2009).
- [21] M.T. Wang, S.T. Yau, *Isometric embeddings into the Minkowski space and new quasi-local mass*, Commun. Math. Phys. 288, 919C942 (2009).
- [22] E. Witten, *A new proof of the positive energy theorem*, Commun. Math. Phys. 80, 381-402 (1981).

- [23] X. Zhang, *A new quasi-local mass and positivity*, Acta Mathematica Sinica, English Series, 24(6), 881-890 (2008).
- [24] X. Zhang, *A quasi-local mass for 2-spheres with negative Gauss curvature*, Science in China Series A: Mathematics, 51, 1644-1650 (2008).

INSTITUTE OF MATHEMATICS, ACADEMY OF MATHEMATICS AND SYSTEMS
SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA

E-mail address: `xzhang@amss.ac.cn`